PROTOCOL FOR NAMING COMPOUNDS IN THE OPCW CENTRAL ANALYTICAL DATABASE

ANY CHANGES TO THIS PROTOCOL SHALL BE APPROVED BY THE VALIDATION GROUP. A NEW VERSION OF THIS PROTOCOL WILL BE DISTRIBUTED TO ALL VALIDATION GROUP MEMBERS.

- 1. In general, the name (spelling, punctuation, spaces, and so on) is to be based on the name given in the Annex on Chemicals to the Chemical Weapons Convention (hereinafter "the Convention").
- 2. The following additional rules should be followed in cases where the information in the Schedules of Chemicals is insufficient to designate only one name.
- 2.1 The name is to be capitalised—the only exceptions being the structural and stereo-descriptors, sec-, tert-, *cis*-, and *trans*-. In cases where a structural or stereo-descriptor prefixes a name, the name is to be capitalised.



2.2 The trivial names for the following radicals are to be used:

Saturated branched Isopropyl, Isobutyl, sec-Butyl, tert-Butyl.



Pinacolyl propylphosphonate 2.B.04

¹ The number listed is the CWC schedule

However, pinacolyl alcohol should be referred to as 3,3-dimethylbutan-2-ol.

Unsaturated Vinyl, Allyl, Isopropenyl.

2.3 When a compound has several substituents, they are to be listed in alphabetical order, irrespective of the presence of N-, O-, or S- prefixes, and of the descriptors, sec-, tert-, cis- or trans-; but see rule 2.5 below.



2.4 The radicals beginning with iso e.g. isobutyl, isopropenyl, and isopropyl are considered to be one entity and are to be listed in alphabetical order starting from "iso" e.g.



2.5 The substituents in Schedule 1.A.03 and 1.B.10 compounds are to be listed in the order "alkyl 2-dialkylaminoethyl", in line with the names given in the Convention, but constituting an exception to rule 2.3 above. The same exception applies to Schedule 2.B.4 compounds containing the "2-dialkylaminoethyl" moiety.



2.6 Parentheses are to be used in the following cases around prefixes defining substituted substituents—after the numerical multiplicative prefixes "bis", "tris", and so on; around simple substituent prefixes to separate locants of the same type referring to different structural elements; and to avoid ambiguity.



- 2.7 For radicals with a branching structure, the name should be derived from the longest continuous chain starting (position 1) at the conjunction with the parent structure. Examples:
 - (a) The methylphosphonofluoridate made using 5-methyl-3-hexanol is 1-Ethyl-3-methylbutyl methylphosphonofluoridate. $F_{0} = P_{1} = 0$ 1-Ethyl-3-methylbutyl methylphosphonofluoridate 1.A.01
 - (b) The name 1-ethyl-2-methylpropyl is to be used instead of 1-isopropylpropyl. $\downarrow 0$ $\downarrow 0$ $\downarrow 0$ $\downarrow 0$ $\downarrow 0$ $\downarrow 1$ -Ethyl-2-methylpropyl ethylphosphonate 2.B.04
 - (c) Substituted chains are to be alphabetical as well.
- 2.8 Thiolate and thionate are to be differentiated according to whether the S-atom is singleor double-bonded to the phosphorus atom.



3,3-Dimethylbutyl S-2-dimethylaminoethyl methylphosphonothiolate 1.A.03



2.9 For phosphorus compounds containing two S-sec-butyl or S-tert-butyl groups linked to phosphorus, the name has to start with Bis(S-sec-butyl) or Bis(S-tert-butyl).



2.10 The name is to be as short as possible, and unnecessary characters such as the following are to be left out



- (b) the 1- before 1-alkyl, in case of a normal alkyl chain;
- (c) the O in O-Alkyl alkylphosphonohalidates;



(d) the O in O-Alkyl S-2-dialkylaminoethyl alkylphosphonothiolates belonging to Schedule 1.A.03; and



Butyl S-2-diisopropylaminoethyl methylphosphonothiolate 1.A.03

- (e) unnecessary brackets and parentheses.
- 2.11 Hydrochloride salts of schedule 2.B.10, 2.B.11 and 2.B.12 chemicals are to be named as free amines with the addition of hydrochloride.

| 2-(N,N-Dipropylamino)ethylchloride hydrochloride | 2.B.10 |
|--|--------|
| 2-(N,N-Dipropylamino)ethanol hydrochloride | 2.B.11 |
| 2-(N,N-Diethylamino)ethanethiol hydrochloride | 2.B.12 |

- 2.12 Substituents to an aromatic ring are to be numbered numerically.
- 2.13 Arsenic-containing derivatives or decomposition products of the Lewisites (Schedule 1.A.05) shall be named as follows:
 - (a) Derivatives with alkyl thioglycolates shall be named per the following examples:
 - (i) 2-Methoxy-2-oxoethyl bis(2-chlorovinyl)arsinothiolite $CI_{As} = S_{O}$
 - (ii) Bis(2-methoxy-2-oxoethyl) 2-chlorovinylarsonodithiolite.



(b) Derivatives with butanethiol shall be named in accordance with the International Union of Pure and Applied Chemistry (IUPAC) rules as follows:



- (ii) Dibutyl 2-chlorovinylarsonodithiolite of the second s
- (iii) Tributyl arsenotrithiolite
- 2.14 Amidate shall be used instead of amidoate (a term used by some commercial chemical naming programs)



Cyclohexyl N,N-diethyl-P-methylphosphonamidate 2.B.04

- 2.15 Compounds with multiple phosphorus atoms shall be named as follows:
 - (a) A symmetrical phosphonate dimer with an oxygen linkage shall be called a pyrophosphonate.



Dicyclobutyl diethylpyrophosphonodithionate 2.B.04

(b) Multiple phosphorus atoms shall be identified as P, P', etc.



4-Methylcyclohexyl N,N-diethyl-P,P'-diethyldiphosphono-P-amidate 2.B.04

(c) Groups attached to the phosphorus shall be indicated by corresponding prime marks (e.g. "-S-methyl" or "-N',N'-diethyl-", etc.), with P listed before P'.



O'-Isopropyl N,N-diethyl-P-methyl-P'-isopropylphosphono-P-amidate 2.B.04

- (d) Otherwise, IUPAC names shall be used.
- 2.16 Compounds based on amines shall be named as follows:
 - (a) Amines named in the Convention shall use the Convention names.
 - (b) Amines with three identical groups shall be named without using the N- prefix. $\overset{c_l}{\overset{c_l}{\overset{}}}$



- (c) All other amines shall be named with the N positions explicitly identified.
- N,N-Diethyl-N-(2-trimethylsilylthioethyl)amine 2.17 Analytical derivatives of amines shall be named on the basis of the original compounds before derivatization.



N-Heptafluorobutyryloxyethyl-N-methyl-N-propylamine DS²



N-Ethyl-N-heptafluorobutyryloxyethyl-N-methylamine DS

2.18 For chemicals other than those listed in the Convention and chemicals with the same configuration around the phosphorus atom, trivalent phosphorus chemicals shall be named according to the IUPAC naming rule that classifies these chemicals as phosphines.



² Derivative of scheduled chemical



2.19 Compounds containing deuterium atoms shall be named as follows: For structure and formula -- use capital D; For names -- use lowercase d.



3. These rules are illustrated below by examples of scheduled compounds, derivatives associated with the scheduled compounds, and non-scheduled reportable chemicals.

| Schedule | Name | Structure |
|----------|---|--|
| 1.A.01 | Alkyl alkylphosphonofluoridate | R |
| | | 0 [≠] [−] \ [°] R' F |
| 1.A.02 | Alkyl N,N-dialkylphosphoramidocyanidate | R AN AN |
| | | |
| 1 \ 03 | Alkyl S 2 dialkylaminoathyl | <u> </u> |
| 1.A.05 | alkylphosphonothiolate | |
| | Alkyl S-trialkylammoniumethyl alkylphosphonothiolate halide (i.e. chloride, iodide) | |
| 1.A.04 | 2-Chloroethylchloromethylsulfide | |
| | Bis(2-chloroethyl)sulfide | |
| | Bis(2-chloroethylthio)methane | CICICI |
| | 1,2-Bis(2-chloroethylthio)ethane | CI S CI |
| | 1,3-Bis(2-chloroethylthio)propane | CI S CI |
| | 1,4-Bis(2-chloroethylthio)butane | CI S CI |
| | 1,5-Bis(2-chloroethylthio)pentane | |
| | Bis(2-chloroethylthiomethyl)ether | |
| | Bis(2-chloroethylthioethyl)ether | |
| 1.A.05 | 2-Chlorovinyldichloroarsine | |
| | Bis(2-chlorovinyl)chloroarsine | |
| | Tris(2-chlorovinyl)arsine | |
| 1.A.06 | Bis(2-chloroethyl)ethylamine | N CI |
| | Bis(2-chloroethyl)methylamine | |
| | Tris(2-chloroethyl)amine | |
| 1.A.07 | Saxitoxin | H ₂ N 0 |
| | | HN NH HN OHOH |
| 1.A.08 | Ricin | |
| 1.B.09 | Alkylphosphonic difluoride | F R—P==0 F |

Examples of Names of Scheduled Compounds

| Schedule | Name | Structure |
|----------|---|---------------------------------------|
| 1.B.10 | Alkyl 2-dialkylaminoethyl alkylphosphonite | N R" |
| | | R—P O—B' |
| 1.B.11 | Isopropyl methylphosphonochloridate | |
| 1.B.12 | Pinacolyl methylphosphonochloridate | CI CI |
| 2.A.01 | O,O-Diethyl S-2-diethylaminoethyl phosphorothiolate | P-S-N |
| 2.A.02 | 1,1,3,3,3-Pentafluoro-2-(trifluoromethyl)-1- propene | F F F F F F F F F F F F F F F F F F F |
| 2.A.03 | 3-Quinuclidinyl benzilate | |
| 2.B.04 | To avoid any confusion, the O and S groups sho sulfur is present. | uld be indicated in esters when |
| | Methylphosphonothionic acid | OH P=S OH |
| | O-Ethyl methylphosphonothionate | |
| | O,O-Diethyl methylphosphonothionate | 0 0-P=S |
| | O-Propyl O-trimethylsilyl propylphosphonothionate | |
| | O-Ethyl S-ethyl methylphosphonothiolate | |
| | S-Ethyl O-methyl methylphosphonothiolate | s |
| | O-Ethyl S-2-methylthioethyl methylphosphonothiolate | |
| | O-Ethyl S-ethyl methylphosphonothiolothionate | |
| | Bis(S-sec-butyl) methylphosphonodithiolate | |

| Schedule | Name | Structure |
|----------|--|-----------------------|
| | O-Ethyl methylphosphonothionochloridate | |
| | S-(4-Mercaptobutyl) | |
| | isopropylphosphonothiolothionochloridate | HS S |
| | Methylphosphonous dichloride | CI P CI |
| | Methylphosphonic dichloride | |
| | Methylphosphonothionic dichloride | CI CI S |
| | Dimethyl methylphosphonate | |
| | Bis(1,2-dimethylpropyl) methylphosphonate | |
| | 1,2-Dimethylpropyl phenyl ethylphosphonate | |
| | Methyl methylphosphonate instead of methyl methylphosphonic acid | 0 |
| | Methylphosphonic acid | OH HO HO |
| | Isobutyl methylphosphonochloridate | |
| | Isopropyl methylphosphonoazidate | |
| | 2-Diisopropylaminoethyl methylphosphinate | D D D H |
| | Methyl 2-diethylaminoethyl methylphosphonate | |
| | O-Ethyl S-2-dibutylaminoethyl methylphosphonothiolate | O THE N |
| | Bis(S-2-diethylaminoethyl) methylphosphonodithiolate | S N O S N |
| | O-Ethyl S-3-diethylaminopropyl methylphosphonothiolate | |
| | Diethyl methylphosphonite | 0-P 0 |

| Schedule | Name | Structure |
|---------------|--|-------------------|
| | Bis(ethylthio)methylphosphine | CH ₃ |
| | | s— |
| | | |
| | | CH ₃ |
| | Dicyclohexyl dimethylpyrophosphonate | |
| | | |
| | | |
| | Dicyclohexyl | |
| | dimethylpyrophosphonodiunonate | |
| | Cyclohexyl N N-diethyl-P- | ŠS A Q (V) |
| | propylphosphonamidate | |
| | | |
| | Isopropyl N,N-diethyl-P,P'- | |
| | dimethyldiphosphono-P-amidate | |
| | O'-Isopropyl N.N-diethyl-P-methyl-P'- | |
| | ethyldiphosphono-P-amidate | |
| | | |
| | N,N,N',N'-Tetramethyl-P,P'- | |
| | dimentyipyrophospholic diamide | |
| | S-Ethyl N,N-dimethyl-P- | P ^P |
| | methylphosphonamidothiolate | ∕_s´ N_ |
| 2.B.05 | N,N-Dialkylphosphoramidic dichloride | R CI |
| | | |
| 2.B.06 | Dialkyl N.N-dialkylphosphoramidate | |
| | | |
| | | |
| | Dimethyl N-ethyl-N-methylphosphoramidate | , <u>0</u> |
| | | N-P=O |
| | | |
| 2.B.07 | Arsenic trichloride | AsCl ₃ |
| 2.B.08 | 2,2-Diphenyl-2-hydroxyacetic acid | HO |
| | | OH OH |
| 2.B.09 | 3-Quinuclidinol | |
| | | |
| 2 D 10 | | / но В |
| 2.B.10 | 2-(IN,N-Dialkylamino)ethylchloride | |
| | 2-(N-Ethyl-N-methylamino)ethylchloride | |
| | | |
| 2.B.11 | 2-(N,N-Dialkylamino)ethanol | R N |
| | | В ОН |

| Schedule | Name | Structure |
|-----------------|--------------------------------------|--------------------------------|
| | 2-(N-Ethyl-N-methylamino)ethanol | N |
| 2 D 12 | 2 (N N Dially lamina) ath an athial | В |
| 2. B .12 | | B-N-SH |
| | 2-(N-Ethyl-N-methylamino)ethanethiol | |
| | | ŃSH |
| 2.B.13 | Bis(2-hydroxyethyl)sulfide | HO |
| 2.B.14 | 3,3-Dimethyl-2-butanol | ОН |
| | | |
| 3.A.01 | Carbonyl dichloride | 0 |
| | | ci Ci |
| 3.A.02 | Cyanogen chloride | CI |
| 3.A.03 | Hydrogen cyanide | H-CN |
| 3.A.04 | Trichloronitromethane | CI NO ₂ |
| | | CI |
| 3.B.05 | Phosphorus oxychloride | CI CI |
| | | P |
| | | |
| 3.B.06 | Phosphorus trichloride | P-Cl ₃ |
| 3.B.07 | Phosphorus pentachloride | P-Cl ₅ |
| 3.B.08 | Trimethyl phosphite | 0 |
| | | |
| 3.B.09 | Triethyl phosphite | Q Q |
| | | |
| 3 R 10 | Dimethyl phosphite | |
| J.D.10 | Difficulty phosphile | |
| | | нограни |
| 3.B.11 | Diethyl phosphite | |
| | | HOPO |
| 3.B.12 | Sulfur monochloride | S ₂ Cl ₂ |
| 3.B.13 | Sulfur dichloride | SCl ₂ |
| 3.B.14 | Thionyl chloride | S(O)Cl ₂ |
| 3.B.15 | Ethyldiethanolamine | |
| 3.B.16 | Methyldiethanolamine | HO |
| 3.B.17 | Triethanolamine | |
| | | , j |
| | | ОН |

Examples of Names for Derivatives of Scheduled Compounds

| Associated Schedule | Chemical Name | Structure |
|------------------------|---|-----------|
| 1.A.05 | 2-(2-Chlorovinyl)-5-methyl-1,3,2- benzodithiarsole | S As Cl |

| Associated Schedule | Chemical Name | Structure |
|------------------------|--|------------------------|
| 2.B.08 | Bis(trimethylsilyl)benzilate | |
| 2.B.09 | 3-Quinuclidinyl trimethylsilyl ether | -O-Si- |
| 2.B.07 | 2-Chloro-5-methyl-1,3,2-benzodithiarsole | S As-Cl |
| 2.B.11 | N,N-Dialkyl-N-(2-trimethylsilyloxyethyl)amine | |
| | N,N-Dialkyl-N-(2-tert- butyldimethylsilyloxyethyl)amine | R N Si A |
| | N-Heptafluorobutyryloxyethyl-N-methyl-N- propylamine | |
| 2.B.12 | N,N-Dialkyl-N-(2-trimethylsilylthioethyl)amine | R N R S Si |
| 2.B.13 | Bis(2-trimethylsilyloxyethyl)sulfide | Si 0 Si |
| 3.B.15 | N-Ethyl-N,N-bis(2- trimethylsilyloxyethyl)amine | |
| 3.B.16 | N-Methyl-N,N-bis(2- trimethylsilyloxyethyl)amine | si |
| 3.B.17 | Tris(2-trimethylsilyloxyethyl)amine | |
| | Tris(2-tert-butyldimethylsilyloxyethyl)amine | |

| Schedule | Chemical Name | Structure |
|----------|--|-----------|
| N.S. | (2-N,N-Diethylaminoethyl)(2-N-ethyl-N- methylaminoethyl)sulfide | N s N |
| N.S. | (2-N,N-Diethylaminoethyl)(2-N-ethyl-N- methylaminoethyl)disulfide | |

Examples of Names for Relevant Non-Scheduled Chemicals